



The *MYC* oncogene — the grand orchestrator of cancer growth and immune evasion

Renumathy Dhanasekaran¹, Anja Deutzmann¹, Wadie D. Mahauad-Fernandez², Aida S. Hansen¹, Arvin M. Gouw² and Dean W. Felsher²✉

Abstract | The *MYC* proto-oncogenes encode a family of transcription factors that are among the most commonly activated oncoproteins in human neoplasias. Indeed, *MYC* aberrations or upregulation of *MYC*-related pathways by alternate mechanisms occur in the vast majority of cancers. *MYC* proteins are master regulators of cellular programmes. Thus, cancers with *MYC* activation elicit many of the hallmarks of cancer required for autonomous neoplastic growth. In preclinical models, *MYC* inactivation can result in sustained tumour regression, a phenomenon that has been attributed to oncogene addiction. Many therapeutic agents that directly target *MYC* are under development; however, to date, their clinical efficacy remains to be demonstrated. In the past few years, studies have demonstrated that *MYC* signalling can enable tumour cells to dysregulate their microenvironment and evade the host immune response. Herein, we discuss how *MYC* pathways not only dictate cancer cell pathophysiology but also suppress the host immune response against that cancer. We also propose that therapies targeting the *MYC* pathway will be key to reversing cancerous growth and restoring antitumour immune responses in patients with *MYC*-driven cancers.

The *MYC* oncogene (also known as *c-MYC*) is part of a superfamily of genes with products that are among the most commonly activated in human cancers^{1–4}. *MYC* is a master regulator of multiple biological programmes and mediates much of its function primarily as a transcription factor that regulates the expression of thousands of genes, either directly or indirectly^{5,6}. In addition, *MYC* exerts multiple biological effects on cellular programmes that influence both the cell-intrinsic biology as well as the host immunity and tumour microenvironment (TME).

MYC has two paralogues: *MYCL* and *MYCN*. The initially held view was that *MYC* is ubiquitously involved in human cancers (both haematological and solid), whereas L-*MYC* is associated with small-cell lung cancer and N-*MYC* with neuroblastoma. However, in the past few years, insights gained from genome-sequencing studies show a broader role for both N-*MYC*^{7–10} and L-*MYC*^{11–13} in many other cancers^{3,4} (FIG. 1a). Thus, when considering the entire *MYC* family, most human cancers harbour the genetic activation of one of its members^{1–4}.

A multitude of reports have documented that *MYC* overexpression can cause tumorigenesis. In a landmark study, Adams et al.¹⁴ demonstrated that transgenic overexpression of *Myc* in mice was sufficient for B cell

lymphomagenesis. Subsequently, many other investigators documented that *MYC* overexpression can causally induce other types of cancer in humans^{15–17}. Moreover, the use of conditional transgenic mouse models has enabled a refined understanding of how *MYC* activation causes tumorigenesis and how its inactivation can elicit tumour regression. Two general approaches have been used to modulate *MYC* in preclinical studies: conditional regulation of the expression of a *MYC* transgene using the *Tet* system^{18–23} and conditional expression of a synthetic *MYC* inhibitor, Omomyc^{24–26}. In these conditional transgenic mouse models, *MYC*-induced tumours regress rapidly and dramatically upon *MYC* inactivation^{18–23}. Tumour regression occurs even without influencing physiological endogenous *MYC* expression or function, in a phenomenon referred to as oncogene addiction²⁷. This phenomenon suggests that targeting *MYC* could be an effective approach to treat some human cancers. To date, however, no drug has demonstrated efficacious therapeutic targeting of *MYC* or the *MYC* pathway, although promising candidates are in development.

Cancer is a complex process that requires a multitude of genetic events and the acquisition of the hallmarks of cancer²⁸. Importantly, *MYC* activation can

¹Division of Gastroenterology and Hepatology, Stanford University School of Medicine, Stanford, CA, USA.

²Division of Oncology, Department of Medicine and Department of Pathology, Stanford University School of Medicine, Stanford, CA, USA.

✉e-mail: dfelsher@stanford.edu
<https://doi.org/10.1038/s41571-021-00549-2>

Key points

- The *MYC* oncogene is activated in the vast majority of cancers by genetic, epigenetic or post-translational mechanisms.
- In preclinical models, inactivation of *MYC* can result in sustained tumour regression owing to oncogene addiction.
- *MYC* activation drives cancer progression through mechanisms involving either the cell-intrinsic acquisition of hallmarks of cancer or dysregulation of the tumour microenvironment and host immune responses.
- *MYC* leads to cancer initiation and maintenance by regulating the host immune system through mechanisms involving immune checkpoints, specific receptors and secreted cytokines.
- Currently, no direct inhibitors of *MYC* are approved; however, many therapeutic agents targeting *MYC* are under development.
- We propose that therapies targeting the *MYC* pathway will be key to reversing cancerous growth and restoring antitumour immune responses in patients with *MYC*-driven cancers.

contribute to many of these hallmarks, including proliferation, self-renewal, cell survival, genomic instability, metabolism and invasiveness as well as angiogenesis and immune evasion^{29–31}. Thus, *MYC* seems to have crucial effects in newly transformed cancer cells and also enables an evolving tumour to remodel the TME and evade the host immune response^{32,33}. Details on how *MYC* biologically modulates cancer cell-intrinsic programmes of cellular proliferation, differentiation, survival and death have been described elsewhere^{34–38}.

In this Review, we focus on the broad clinical importance of *MYC* and *MYC*-related oncogenic signalling in the pathogenesis of human cancers. In particular, we provide evidence that *MYC* is a major driver of human cancer through its effects on tumour cells and also by enabling cancers to evade host immune surveillance. We summarize the studies that have demonstrated rapid and sustained tumour regression in *MYC*-driven cancers upon *MYC* inactivation as a consequence of oncogene addiction. Finally, we provide a rationale for and examples of therapeutic strategies targeting the *MYC* pathway. We conclude that *MYC* signalling is one of the most important, yet to be successfully targeted, oncogenic pathways in human cancer; thus, mechanistic insights into the tumorigenic roles of this gene family will be key to developing efficacious therapies for cancer.

***MYC* is often activated in human cancers**

Genetic alterations affecting the *MYC* proto-oncogenes and the *MYC*-related signalling pathways are among the most common in human cancers^{1–4} (FIG. 1). *MYC* can be activated in cancers through multiple genetic, epistatic, epigenetic and post-translational mechanisms (FIG. 2), which vary between cancer types.

Genomic alterations, including gene amplification, chromosomal translocations and mutations, can increase *MYC* expression^{1,4,39} (FIG. 2a). A pan-cancer assessment of alterations in 33 human cancer types conducted as part of The Cancer Genome Atlas project revealed that *MYC* or its paralogues *MYCL* or *MYCN* are amplified in 28% of tumours³. An analysis encompassing 17 cancer types also showed that *MYC* is frequently amplified in extrachromosomal DNA during tumour evolution^{40,41}. These amplifications can directly lead to *MYC*

overexpression or, indirectly, to the activation of genes involved in the *MYC* pathway as demonstrated in 30 of the 33 cancer types analysed in The Cancer Genome Atlas pan-cancer study³. *MYC* is genetically amplified in many solid tumour types, including breast and liver cancers, and is frequently chromosomally translocated in B cell and T cell leukaemias and lymphomas^{39,42,43} (FIG. 1). *MYC* expression can also be increased through the insertion of upstream enhancers by retroviruses⁴⁴ or epistatically and/or epigenetically through the activation of many other oncogenic pathways, such as those mediated by WNT- β -catenin, SRC, numerous receptor tyrosine kinases and Notch⁴⁵, or via the loss of tumour suppressors such as APC⁴⁶ and TGF β ⁴⁷ (FIG. 2b).

In addition, *MYC* can be activated post-translationally through many mechanisms increasing protein stability^{48,49} (FIG. 2c). *MYC* has a short half-life that is tightly regulated through phosphorylation and proteasomal degradation. Tumours with stable *MYC* expression have elevated levels of phospho-serine 62 (P-S62)-*MYC* and low levels of phospho-threonine 58 (P-T58)-*MYC*^{49,50}. Mitogenic pathways, such as RAS-MEK-ERK signalling, can increase P-S62 levels and thereby increase *MYC* stability⁵¹. Moreover, *MYC* mutations affecting the T58 residue can lead to constitutive S62 phosphorylation⁵². Cancers can also down-regulate PP2A, a serine/threonine phosphatase complex that targets P-S62, leading to *MYC* accumulation^{50,53–55}. The levels of PIN1, a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family that functionally regulates *MYC* stability via isomerization, is overexpressed in multiple human cancer types, suggesting that this is another mechanism by which levels of *MYC* are increased in cancer cells^{56,57}. Finally, FBW7 is a tumour suppressor that controls the proteasomal degradation of many proteins, including *MYC*⁵⁸. FBW7 is frequently inactivated in human cancers, such as uterine (18%), colon (16%) or cervical cancer (13%), through deletion, mutation or epigenetic modification^{59,60}, which can promote cancer progression at least in part owing to the resulting increase in *MYC* levels⁵³.

The *MYC* superfamily also includes genes encoding multiple other proteins considered to function as transcription factors or co-regulatory proteins and which are very commonly activated in human cancers⁶¹. These genes include *MAX*, which encodes an *MYC* heterodimer binding partner, and *MLX* as well as *MGA*, *MXD1*, *MXD3*, *MXD4*, *MXI1*, *MNT*, *MLXIP* and *MLXIPL*, which encode proteins that interact with *MAX* and/or *MLX*⁶¹. Interactions between different members of this superfamily can result in transcriptional activation or repression of specific genes that, in turn, regulate cell-cycle progression and/or cellular transformation. Mutations or deletions in these members of the *MYC* superfamily have also been reported in human cancers^{62,63}. Therefore, the *MYC* pathway is dysregulated in the majority of human cancers^{3,29,64,65}. Accordingly, we have suggested that *MYC* signalling is a molecular hallmark of cancer⁶⁶.

MYC activation alone is generally not sufficient to induce the neoplastic transformation of non-malignant cells. Indeed, the tumorigenic functions of oncogenic

MYC are restrained by many physiological mechanisms that cause cell-cycle arrest, apoptosis and/or cellular senescence. For example, unrestricted MYC activation can induce the expression of TP53 (REF.⁶⁷) as well as of cell-cycle checkpoint genes (such as CDKN2A)⁶⁸ and/or regulators of apoptosis (such as BCL2)⁶⁹. Correspondingly, the inactivation of genes such as TP53 or CDKN2A, the activation of genes such as BCL2, or the deletion of pro-apoptotic genes (for example, that encoding caspase-8 (REF.⁷⁰)) cooperate with MYC signalling to block cell-cycle arrest, senescence and/or apoptosis, thereby driving cancer initiation. The progressive shortening of the telomeres after every cell cycle eventually halts cell division and is another cellular protective mechanism; however, MYC increases telomerase activity by inducing the expression of a telomerase reverse transcriptase, thus favouring cellular immortality⁷¹.

Provocatively, despite activation of MYC signalling being one of the most common genetic events in human tumorigenesis, MYC might not always be essential for tumour initiation. Genetically engineered

Myc-haploinsufficient (*Myc*^{+/-}) mice develop lymphomas at the same rate as *Myc*-wild-type mice, albeit with reduced cell proliferation and delayed tumour progression⁷². Regardless, members of the MYC superfamily are commonly activated and thought to be required for tumorigenesis in most human cancers.

MYC oncogene addiction

In 2002, Bernard Weinstein proposed that cancers are ‘oncogene addicted’²⁷ following evidence from both experimental mouse models of MYC-driven cancers and clinical studies of heterogeneous human tumours. A multitude of studies has demonstrated that MYC can act as a cancer driver and that suppression of MYC signalling can result in sustained tumour regression^{18–23}. In transgenic mouse models of MYC-driven cancers (including lymphoma, leukaemia, osteosarcoma, hepato cellular carcinoma (HCC), renal cell carcinoma and lung adenocarcinoma (LAC)), sustained tumour regression can be observed upon MYC inactivation, although the specific consequences depend upon the particular type of cancer^{18–23}. Furthermore, the reversible expression of a dominant-negative MYC mutant and MYC-MAX disruptor, Omomyc, leads to the rapid regression of lung tumours in mouse models^{24–26}. Thus, extensive experimental evidence supports targeting MYC as a potential therapeutic approach against a multitude of human cancers. MYC activation drives cancer progression through two types of mechanisms involving either acquisition of hallmarks of cancer that are intrinsic to tumour cells^{2,66} or changes in the TME and anticancer immune response^{32,73}.

Effects of MYC activation

Cancer cell-intrinsic mechanisms. MYC regulates cancer cell-intrinsic processes with several physiological roles such as cell growth, differentiation and metabolism^{74–77}. Such processes are co-opted upon MYC activation in a cancer cell-autonomous manner, thus facilitating cellular transformation (FIG. 3). In non-malignant cells, the expression of MYC is tightly controlled. High levels of MYC expression are, in fact, associated with increased sensitivity to apoptosis^{78,79}. However, MYC activation can overcome these physiological barriers and promote cancer through several interdependent cell-intrinsic mechanisms.

MYC overexpression enforces relentless cellular proliferation by enabling cancer cells to re-enter the cell cycle⁷⁹ and also accelerates progression through the cell cycle by inhibiting cell-cycle checkpoints^{37,80,81}. In addition, MYC promotes cancer cell growth by inducing a substantial increase in ribosomal and protein biogenesis^{82–85}. MYC alters cancer cell metabolism by globally rewiring multiple metabolic pathways to support rapid growth and proliferation^{35,86}. This function is partially accomplished by facilitating nutrient uptake in cancer cells through transcriptional induction of glucose and glutamine transporters and also of genes involved in glycolysis and glutaminolysis^{87–89}. MYC activates cellular survival programmes through specific effects on DNA replication. In particular, MYC can directly activate the DNA replication machinery⁹⁰. Of note, *Myc* is one






Fig. 1 | Major genetic alterations involving MYC and its paralogues in human cancers. Prevalence of gene amplification of the three MYC paralogues MYC, MYCL and MYCN across 16 major human cancer types in The Cancer Genome Atlas.

a Genetic aberrations



b Transcriptional regulation



c Protein stabilization




Fig. 2 | Mechanisms leading to MYC activation in human cancers. a Genetic aberrations, such as chromosomal translocations and genomic amplifications, lead to increased MYC mRNA expression. **b** Alteration of upstream regulatory pathways can lead to increased or decreased transcription of the MYC oncogene. **c** Post-translational modifications of the MYC protein, such as preferential phosphorylation of the serine 62 (S62) residue versus threonine 58 (T58), can block degradation and promote stabilization of MYC, thereby enhancing activation of the MYC pathway.

of the four factors that enable stemness and embryonic programmes of self-renewal in induced pluripotent stem cells⁹¹. In addition, MYC can bypass and/or transcriptionally repress cell-cycle checkpoint proteins, such as p15 or p21, in some contexts, thus preventing cellular senescence even in the presence of DNA damage⁹². MYC activation can also result in genomic instability owing to many types of genomic damage, including DNA breaks, chromosomal translocations, chromosomal gains or losses, aneuploidy, and polyploidy. The mechanisms by which MYC induces genomic damage involve overriding cell-cycle checkpoints (such as transcriptional repression of p21) and blocking DNA double-strand break repair^{21,93–96} as well as mitochondrial reactive oxygen species generated during MYC-driven cell proliferation^{97–99}. MYC activation also can induce proliferative arrest and endoreduplication as well as genomic instability *in vivo*^{100,101}.

The ability of oncogenic MYC to elicit cell proliferation seems to depend upon the specific tissue lineage and the developmental context. For example, MYC overexpression in liver cells of embryonic or neonatal mice results in rapid cellular proliferation and rapid formation of hepatoblastomas (a paediatric neoplasm), whereas MYC overexpression in liver cells of adult mice results in cellular hypertrophy, endoreduplication and late-onset HCC¹⁰⁰. In summary, MYC activation contributes to both the initiation and maintenance of tumorigenesis through several cancer cell-intrinsic mechanisms.

Effects on the immune response and TME. Preclinical studies have revealed many mechanisms by which MYC influences the TME, involving effects on host stromal cells, vascular endothelial cells, innate and adaptive

immune cells, and, in some cases, specific cytokines and their receptors (FIG. 4).

MYC expression in cancer cells can influence host endothelial cells, resulting in reprogramming of the TME to support cellular proliferation and induction of angiogenesis^{102–104} (FIG. 3). Moreover, MYC initiates and maintains tumorigenesis through a thrombospondin 1 (TSP1)-regulated angiogenic switch¹⁰⁵ (FIG. 3).

MYC-dependent perturbation of the host immune response is causally related to the mechanism of tumorigenesis and is an important mechanism underlying tumour regression upon MYC inactivation^{106–108}. Characterized mechanisms include effects on CD4⁺ T cells, macrophages, and natural killer (NK) cells and involve cytokines (such as TGFβ, interferons and TSP1) and key immunomodulatory molecules (such as PD-L1, CD47 and MHC class I (MHC I); FIG. 4)^{106–112}. More than 30 years ago, a study described that MYC downregulates the expression of proteins involved in antigen presentation, such as MHC I, thus enabling immune evasion^{113–115}. In addition, MYC induces the secretion of immune inhibitory cytokines from cancer cells, such as TGFβ, which in turn suppresses cytolytic T cell responses^{116,117}. Furthermore, MYC has been found to upregulate the expression of immune-checkpoint proteins, such as PD-L1 and CD47, thus leading to T cell exhaustion^{107,108,110,118,119}. In addition, MYC suppresses NK cell-mediated immune surveillance by transcriptionally repressing STAT1 and STAT2 and the type I interferon pathway^{107,109,111}. Finally, MYC cooperates with TWIST1 to regulate the secretion of CCL2 and IL-13, thereby promoting the recruitment and polarization of immunosuppressive macrophages, which in turn facilitate metastasis¹²⁰ (FIG. 4).

Effects of MYC inactivation

MYC inactivation can cause tumour regression through effects on tumour cells as well as the host immune system and TME. In preclinical models^{18–23}, MYC inactivation results in a cascade of changes in both tumour cells and the TME that lead to rapid tumour regression and the restoration of regular tissue structures (FIG. 4). The specific kinetics and consequences of MYC inactivation seem to depend upon the tissue of origin of a cancer as well as on the specific genetic context. Thus, in mice with MYC-driven haematological cancers, such as T cell acute lymphoblastic leukaemia, MYC inactivation is associated with rapid proliferative arrest and very rapid tumour regression within a few days, together with robust induction of cellular differentiation, senescence and apoptosis²⁰. By contrast, in mice with MYC-driven mesenchymal-derived tumours, such as osteosarcoma, MYC inactivation results in the robust terminal differentiation of bone cancer into bone^{18,21}. Finally, in mice with

MYC-driven epithelial-derived tumours, such as HCC, LAC and renal cell carcinoma, MYC inactivation results in tumour regression but a dormant population of otherwise histologically normal-appearing cells remains^{19,121,122}.

Cancer cell-intrinsic mechanisms. MYC inactivation leads to the restoration of physiological cell-intrinsic checkpoints on cellular growth and differentiation. In mouse models of MYC-driven cancer, tumour cells undergo proliferative arrest, differentiation, senescence and/or apoptosis upon MYC inactivation^{18–23,66}. These processes seem to occur through the restoration of cell-cycle checkpoints, DNA repair and chromatin remodelling programmes. Thus, the knockdown of cell-cycle checkpoint proteins⁶⁶, such as p16, or DNA repair regulators, such as p53, impedes tumour regression elicited by MYC inactivation²¹. MYC inactivation also causes tumour regression through the loss of survival signals with the maintenance of death signals within cancer





Fig. 3 | MYC is a grand orchestrator of the hallmarks of cancer. MYC regulates several cancer cell-intrinsic and host-dependent pathways to promote cancer cell growth and survival (green area). Cancer cell-intrinsic processes regulated by MYC include proliferation, metabolism, invasiveness, autophagy, and protein and ribosomal biosynthesis. MYC also simultaneously blocks other cellular protective mechanisms, such as differentiation or senescence, thereby promoting cancer progression (red area). Paradoxically, MYC also induces cellular processes, such as apoptosis and chromosomal instability, that can be detrimental to cancer cell survival (orange arrows). The delicate balance between these events and cellular context ultimately determines cell fate. Furthermore, MYC controls the ability of cancer cells to undergo dormancy and to overcome nutrient-low environments, eventually leading to tumour relapse. To maintain this quiescent state, MYC inhibits several cellular programmes, including cell differentiation and senescence. MYC activation in the cancer cells also drives enhanced angiogenesis, thus promoting tumour progression. Finally, one of the most important functions of MYC is to enable cancer cells to evade and inhibit immune surveillance to safeguard their survival. All of these hallmarks regulated by MYC work in unison to drive cancer progression.

a MYC-ON



b MYC-OFF




Fig. 4 | MYC blocks immune surveillance. MYC enables cancers to evade the immune system through several distinct mechanisms. **a** | MYC regulates the expression and production of several immune ligands or receptors and immune effector molecules, such as PD-L1, CD47, MHC classes I and II, and NKG2D. MYC also promotes the expression of several cytokines, such as CCL2, IL-23 and CCL9, which regulate the conversion of antitumour M1 macrophages to pro-tumour M2 macrophages and prevent the activation and recruitment of B cells, natural killer (NK) cells, and CD4⁺ and CD8⁺ T cells. CCL9 activates mast cells, which in turn induce angiogenesis. **b** | Upon inactivation of MYC, the downregulation of PD-L1 and CD47

results in the rapid recruitment and activation of CD8⁺ T cells and NK cells. The inactivation of MYC also increases the levels of NKG2DL in cancer cells, resulting in NK cell recruitment. The production of most of the cytokines described above decreases upon MYC inactivation. By contrast, the expression of type I interferons and CCL5 increases upon MYC inactivation, resulting in the recruitment and activation of NK cells and B cells and of CD8⁺ T cells, respectively. Thus, MYC controls the immune status of a tumour by creating an immunosuppressive ‘cold’ tumour microenvironment when activated, which reverts to an immune-sensitive ‘hot’ milieu when inactivated. TSP1, thrombospondin 1.

cells¹²³. The mechanism involves both a TGFβ autocrine mechanism that elicits proliferative arrest and senescence upon MYC inactivation^{116,123} and the modulation of microRNAs (such as miR17-92, miR-15 and let-7 family members) that epigenetically regulate genes required for cell survival and stemness^{124–127}. Thus, MYC maintains tumorigenesis partially through the coordination of multiple cell-intrinsic biological programmes.

Effects on the immune response and TME. MYC inactivation can also elicit tumour regression through effects on the host immune response and TME such as the suppression of angiogenesis^{102,105}. Indeed, MYC has a known role in inducing the expression of cytokines that regulate angiogenesis. One such key regulator is the anti-angiogenic factor TSP1, which is transcriptionally repressed by MYC¹²⁸. Upon MYC inactivation, the upregulation of TSP1 results in the death of endothelial cells within the tumour vasculature and the loss of mean vessel density¹⁰⁵.

Within a week of MYC inactivation, host innate immune cells (such as macrophages^{107,120} and NK cells^{111,129}) and adaptive immune effectors (including CD4⁺ T cells^{106,130} and B cells²³) are recruited to the tumour bed (FIG. 4). All of these changes in host cellular compartments have been correlated and, in some cases, shown to be causally associated with changes in the levels of specific cytokines (such as CCL2 or IL-1β) and immune checkpoints (for example, PD-L1 or CD47)^{103,107,108,120}. After several weeks of MYC inactivation, complete regression of the tumour occurs together with the restoration of regular tissue architecture and establishment of a durable immune response. Indeed, tumour regression upon MYC inactivation seems to be associated with long-standing immune rejection of the same tumour type¹⁰⁶.

Modelling oncogene addiction. Of note, changes associated with oncogene addiction can be modelled mathematically. For example, one such model was developed to

predict how relative changes in survival and death signals affect tumour growth¹²³. In transgenic mouse models of LAC driven by *MYC* alone, *Ras* alone or both, oncogene addiction was found to be best described as a consequence of a reduced proliferative response rather than a net increase in apoptosis¹²³. This observation is consistent with the ‘oncogenic shock model’, which proposes that, upon inactivation of oncogenic pathways, pro-survival signals dissipate quickly whereas pro-apoptotic signals persist for longer^{123,131}. Indeed, the genetic context of a tumour and its sensitivity to undergo proliferative arrest and/or apoptosis influences the ability of *MYC* inactivation to elicit tumour regression. The inactivation of oncogenes other than *MYC*, such as *RAS* or *ERBB2*, can lead to similar phenotypical features of oncogene addiction, suggesting that the mechanisms involved are not necessarily unique to *MYC*^{132–134}. Lastly, mathematical models have demonstrated mechanisms by which the TME contributes to tumour regression upon oncogene inactivation in transgenic mouse models¹³⁵.

Mechanisms of host immune evasion

MYC activation in cancer cells can impede the host immune response. Conversely, its inactivation seems to restore antitumour immunity in a sequential manner. In an experimental mouse model of pancreatic cancer, *MYC* activation led to the rapid induction of the potent pro-inflammatory cytokine IL-1 β , driving the increased proliferation of endothelial cells¹⁰³. *CCL2* and *CCL5* were also induced, thus triggering the recruitment of mast cells and delaying that of macrophages and neutrophils¹³⁶. In 2020, Muthalagu et al. reported a novel mechanism whereby *MYC* and *KRAS-G12D* cooperatively promote pancreatic cancer progression in mice by repressing the type I interferon pathway¹¹¹. Targeted suppression of *MYC*–*MIZ1* complex formation restores the activation of this pathway and leads to *CXCL13*-mediated recruitment of antitumour B cells and NK cells¹¹¹. Therefore, in multiple mouse models, *MYC* inactivation is associated with the restoration of an anticancer innate and adaptive immune response.

Some studies have revealed that *MYC* can regulate the immune response in a manner clearly dependent upon another driver oncogene. For example, in an experimental mouse model of HCC, *MYC* activation in the context of *TWIST1* activation leads to transcriptional changes that confer macrophages from a pro-inflammatory to a prometastatic phenotype mediated by the secretion of *CCL2* and IL-13 (REF.¹²⁰). Importantly, levels of *MYC*, *TWIST1*, *CCL2* and IL-13 were also directly correlated with a worse prognosis in 33 different human cancers. Moreover, measurement of *CCL2* and IL-13 enabled prediction of the invasiveness of HCC in patients with this cancer type. Similarly, *MYC* and *RAS* cooperate to elicit an inflammatory phenotype in a mouse model of LAC¹⁰⁷ through a process characterized by increased production of *CCL9* and IL-23 (REF.¹⁰⁷). In this model, PD-L1⁺ macrophages inhibit T cells and induce angiogenesis. In summary, the specific features of the immune response regulated by *MYC* seem to be dependent on the specific type of cancer as well as on the genetic events associated with that tumour¹³⁷.

MYC regulates immune checkpoints

MYC affects the host immune response through the regulation of multiple immune checkpoints, including PD-L1 and CD47. *MYC* regulates PD-L1 in a complex manner involving multiple mechanisms, including direct transcriptional regulation as well as indirect post-transcriptional regulation^{108,138,139}. In transgenic mouse models of haematological cancers, *MYC* binds to the promoter region and regulates the transcription of *Cd274* (encoding PD-L1)¹⁰⁸. The constitutive retrovirally mediated expression of PD-L1 abrogated the ability of *MYC* inactivation to result in the inhibition of angiogenesis and the induction of cellular senescence and blocked tumour regression^{108,72,107,140,141}. Similarly, in neuroblastoma and *ALK*-negative anaplastic large cell lymphoma cells, the in vitro inhibition of *MYC* expression with short-hairpin RNAs or small-molecule inhibitors reduces PD-L1 expression^{142,143}. Thus, the transcriptional regulation of *CD274* expression seems to be a conserved mechanism by which *MYC* promotes immune evasion.

MYC can also regulate PD-L1 indirectly in cooperation with other oncogenes or through other key transcriptional regulators. For example, in a mouse model of triple-negative breast cancer, *MYC* and p65 cooperate to regulate *Cd274* transcription¹¹⁰. In mice with LAC, *MYC* cooperates with *KRAS* to overcome translational repression of PD-L1 (REF.¹⁰⁷). Increased protein stability of *MYC* is another mechanism of PD-L1 regulation. Indeed, in a transgenic mouse model of HCC, an engineered increase in *MYC* protein stability correspondingly increased PD-L1 expression and reduced the tumour infiltration of CD8⁺ T cells¹¹⁸. In summary, *MYC* can increase the expression of PD-L1 in cancer cells through multiple mechanisms.

MYC can also regulate other receptors involved in the immune response such as CD47 and MHC I. CD47 is an immune regulatory molecule that provides a ‘do not eat me’ signal to immune cells expressing two types of ligands: TSP1 and tyrosine-protein phosphatase non-receptor type substrate 1 (also known as SIRP α)^{144,145}. In human-derived lymphoma cells, *MYC* upregulated CD47 and inhibition of *MYC* signalling with the BET bromodomain inhibitor JQ1 reduced CD47 levels without affecting PD-L1 expression¹⁴⁶. Similarly, in a mouse model of cutaneous T cell lymphoma, high levels of *MYC* induced the transcription of *CD47* (REF.¹⁴⁷). Interestingly, CD47 has been shown to increase *MYC* expression in a feedforward loop^{144,147}.

Finally, *MYC* suppresses the expression of MHC I. More than 30 years ago, *MYC* was shown to downregulate MHC I in multiple cancers, including neuroblastoma, melanoma and lymphoma^{113,115,148}. A subsequent study in mice showed that this downregulation is associated with a decrease in the recruitment of T cells and NK cells to LACs¹⁰⁷. Of note, oncogenic *RAS* also seems to decrease MHC I expression; this process might occur through its known cooperation with *MYC*^{149,150}. Together, these studies show that the regulation of immune checkpoints is a crucial mechanism whereby the *MYC* oncogene remodels the immune TME and facilitates immune evasion.

Immune regulation through metabolism

MYC regulates diverse metabolic pathways in cancer cells^{151–154}. Importantly, immune function is also highly regulated by cellular metabolism^{155–158}. More specifically, MYC influences the host TME directly and indirectly through its effects on metabolism.

MYC expression in cancer cells results in perturbations in metabolic pathways that lead to the release of immunomodulatory molecules. During tumour initiation in mouse models, MYC-driven proliferating cancer cells release metabolites (such as lactate¹⁵⁹ and glutamate^{160,161}) that influence the local immune function within the TME. As the tumour grows, these effects could become more general and cause systemic disruption of anticancer immune responses^{162,163}.

MYC-driven cancers have dysregulated glucose, glutamine and lipid metabolism, which could affect the host metabolic homeostasis thereby indirectly influencing the immune system and immunoediting^{152,164–166}. MYC enables a metabolic shift from oxidative phosphorylation to glycolysis (the Warburg effect), upregulates glutaminolysis and increases lipogenesis in cancer cells^{151–154}, which improves cellular fitness and confers a cell survival advantage. However, such an advantage could lead to a competition between cancer and immune cells for key metabolites, thus making immune cells ineffective¹⁶⁷. Hence, MYC could have both direct and indirect effects on the host immune system through its effects on cancer metabolism.

The MYC pathway as a therapeutic target

Despite substantial experimental evidence showing that targeting MYC can lead to tumour regression through both direct and indirect effects^{18–26} and decades of effort towards clinical translation of these findings, MYC has not yet been successfully targeted therapeutically. The difficulties in targeting MYC have been described elsewhere^{168,169}; briefly, some of the reasons include the highly disordered protein structure of MYC and the lack of a binding pocket or specific enzymatic activity. Nevertheless, a variety of strategies to inhibit MYC activity are currently being explored (FIG. 5) in preclinical cancer models or in clinical studies (Supplementary Table 1). Individual approaches have been comprehensively reviewed elsewhere^{169–175}; herein, we provide a broad overview of different strategies for the therapeutic targeting of MYC-driven cancers.

Given that MYC functions as a heterodimer with MAX¹⁷⁶, small molecules that target the MYC–MAX interface, stabilize MAX–MAX homodimers or disrupt the binding of MYC–MAX to DNA have been used to inhibit MYC signalling. The MYC–MAX disruptors MYCi361 and MYCi975 (REF.¹⁷⁷), the MAX–MAX stabilizer KI-MS2-008 (REF.¹⁷⁸), disruptors of MYC–MAX binding to its canonical Ebox DNA sequence (such as ME47 (REF.¹⁷⁹)), the inhibitor of MYC–DNA binding and MYC transcriptional activity EN4 (REF.¹⁸⁰), and Omomyc have all shown activity in preclinical models of Myc-driven cancers^{24,25}. Some of the discovered small molecules and designed miniproteins show efficacy in various preclinical mouse cancer models but clinical evidence is not yet available. To date,

Omomyc is the only promising candidate to have entered clinical trials.

Another approach to targeting MYC involves decreasing MYC biosynthesis or altering its stability. For example, inhibitors of the PI3K–AKT–mTOR signalling pathway suppress MYC translation and thus decrease tumoural levels of MYC in a mouse model¹⁸¹. Similarly, silvestrol, a small-molecule inhibitor of the translation initiator eIF4A, reduces MYC translation and inhibits tumour growth in a mouse model of colon cancer¹⁸². Moreover, inhibitors of Aurora kinases can induce MYC protein degradation and specifically inhibit MYC protein overexpression in cancer cells without affecting physiological MYC expression in non-malignant cells^{183,184}. PLK1 also regulates MYC protein stability and is another promising therapeutic target¹⁸⁵. Finally, PIN1 modulates MYC turnover and transcriptional activity¹⁸⁶. Specific PIN1 inhibitors have shown promising therapeutic activity in preclinical models of MYC-driven cancers¹⁸⁷.

In the past few years, important developments have occurred in the design of specific protein degraders or proteolysis-targeting chimaeras (PROTACs). These approaches are based on cereblon-mediated protein degradation and provide a new means to directly and specifically target transcription factors, such as MYC and/or its interaction partners, for proteasomal degradation^{188,189}.

Reducing MYC mRNA stability in tumours is another potential strategy to target MYC-driven tumours¹⁹⁰. Antisense oligonucleotides have been used in preclinical models to target MYC^{191–193}. For example, a MYC-specific antisense oligonucleotide, MYC-ASO, impairs tumour progression and elicits an antitumour immune response in a primary transgenic mouse model of HCC¹⁹⁴.

Another strategy involves targeting upstream regulators to inhibit MYC transcription. For example, BET-motif inhibitors have demonstrated antitumour efficacy that might partially result from the inhibition of *Myc* transcription^{195,196}. Drugs that stabilize the G-quadruplex structure in the *Myc* promoter region, such as IZCZ-3, can also inhibit *Myc* transcription^{197–201}.

Vulnerabilities that might be therapeutically targetable can be identified in screens for MYC-related synthetic lethal interactions — that is, of genes likely to be key mediators of MYC-driven cellular processes required for tumorigenesis. As a transcription factor, MYC drives the expression of a multitude of gene products that are required to initiate and/or maintain cancers^{2,5,61,202,203}, thus diversifying the arsenal of potential drug targets. Synthetic lethal screens^{204–206} have identified multiple potential targets, such as cyclin-dependent kinases (CDKs)^{207,208}, SAE1/2-mediated sumoylation²⁰⁴ and TNFRSF10B²⁰⁵. In addition, studies have demonstrated the dependency of MYC-driven cancers on a functional DNA damage response mediated by ATR, CHK1 and PRKDC^{209,210}. A novel therapeutic synthetic lethal approach could address the transcription-independent role of MYCN in preventing replication stress-induced DNA damage. Indeed, in complex with Aurora kinase A, MYCN prevents R-loop formation and thus facilitates cell proliferation in a mouse model of neuroblastoma. This process could potentially be exploited therapeutically through the combined inhibition of Aurora kinase A and ATR²¹¹.




Fig. 5 | **Therapeutic strategies to target MYC-driven tumours.** Among the multiple strategies currently explored to target MYC-driven tumours, the majority use indirect approaches (grey boxes) such as those based on inhibiting MYC synthetic lethal genes or interfering with the expression of MYC at the DNA, RNA or protein level. Direct strategies to inhibit MYC (blue box) include approaches using small molecules, peptides or ‘miniproteins’ to inhibit MYC–MAX dimerization, sequester MAX via homodimer stabilization, or interfering with MYC–MAX binding to target DNA sequences. Ac, acetylation; CDKs, cyclin-dependent kinases; HDACs, histone deacetylases; Me, methylation; P, phosphorylation; Ub, ubiquitylation.

In *Myc*-driven mouse models of B cell lymphomas or lung cancer, the anti-apoptotic protein MCL1 is a synthetic lethal vulnerability of MYCN. MCL1 inhibitors are currently being tested in clinical trials (Supplementary Table 1)^{212,213}. Of note, these mechanisms are specific to MYCN.

Epigenetic modifiers, such as histone deacetylases (HDACs)^{214–216} or histone methyltransferases (such as EZH2²¹⁷), can also be used to target *MYC*-driven cancers. In non-small-cell lung cancer cell lines, the combination of DNA methyltransferase inhibitors with HDAC inhibitors led to the suppression of MYC signalling and consequent activation of an IFN α / β -based transcriptional programme accompanied by a CCL5-mediated antitumour T cell response¹⁴⁰. These investigators were able to demonstrate that epigenetic therapy could deplete MYC, enabling tumours to reverse immune evasion¹⁴⁰. Similarly, MYC can promote cancer through miR17-92-mediated regulation of chromatin remodelling^{104,125,218}. The use of anti-miR-17 oligonucleotides delayed

tumour progression in a mouse model of liver cancer²¹⁹. Finally, given that MYC is a key regulator of cellular metabolism^{19,35,152}, targeting MYC synthetic metabolic vulnerabilities, such as SREBP1, is another potential therapeutic strategy^{153,165}. Our laboratory is currently exploiting the concept of synthetic lethality with CRISPR-based functional in vivo genomic screening to identify novel molecular targets in *MYC*-driven cancers (A.D. and D.W.F., unpublished results).

A final therapeutic approach involves exploiting MYC-induced alterations of immune surveillance. For example, MYC regulates the expression of specific immunomodulatory molecules and cellular immune mechanisms in specific cancers and, thus, molecular and cellular therapies that restore these mechanisms might be particularly effective against *MYC*-driven cancers. Some evidence supports this strategy, at least experimentally, such that *Myc*-driven mouse cancers have been shown to be sensitive to the cytokine-mediated

restoration of anticancer immune responses²²⁰, reinstatement of NK cell-mediated immune surveillance¹⁰⁹ or immune-checkpoint inhibition^{31,108}.

Similarly, drugs that target the MYC pathway might sensitize tumours to specific immunotherapies. For example, MYCi361 has demonstrated promising synergy with anti-PD-1 antibodies in a mouse model of *Myc*-driven prostate cancer¹⁷⁷. In addition, in a xenograft model of pancreatic ductal adenocarcinoma, the combination of an anti-PD-L1 antibody and JQ1 had synergistic effects²²¹. Hence, MYC expression might predict response to immunotherapy and targeting the MYC pathway might sensitize cancers to and enhance the efficacy of immunotherapy.

In summary, many strategies can be used to either directly or indirectly target *MYC*-driven cancers. Ongoing early phase clinical trials are testing direct MYC inhibitors that either silence *MYC* gene expression, inhibit MYC protein biosynthesis or target MYC for proteasomal degradation (Supplementary Table 1). MYC can also be inhibited indirectly by targeting identified synthetic lethal interactions, with examples including HDACs, CDK4, CDK6, CDK7 and CDK9, DNA repair genes (such as CHK1 and ATR), and anti-apoptotic genes (for example, MCL1; Supplementary Table 1). Furthermore, small-molecule screens have identified compounds that directly target MYC–MAX. Nevertheless, most of these MYC–MAX inhibitors are yet to enter clinical trials. Strategies such as antisense RNA oligonucleotides have been tested in phase I clinical trials but, to our knowledge, have not been further pursued as cancer therapeutics^{222,223}. To date, Omomyc is the only agent directly targeting MYC that has been tested in clinical trials²²⁴ (Supplementary Table 1).

We believe that direct targeting of MYC is likely to be more clinically effective than indirect targeting strategies. Nevertheless, several drugs targeting MYC indirectly, either through synthetic lethal interactions or upstream signalling pathways, have reached the clinic (Supplementary Table 1). These agents include CDK4/6 inhibitors and mTOR inhibitors; however, such approaches are likely to be associated with therapeutic resistance. As we have described, MYC controls interconnected cell-intrinsic and cell-extrinsic programmes and, thus, cancer cells will probably be able to bypass the inhibition of any one specific mechanism. Therefore, even the brief or partial depletion of MYC can result in sustained suppression of tumorigenesis in preclinical models^{18,105,225}. We believe that future efforts will make it possible to find agents that directly target MYC and that a therapeutic window exists in which these drugs have clinical activity against cancer while mitigating toxic effects in non-malignant cells.

Future directions

Insights on how MYC regulates both cancer cell-intrinsic processes, such as proliferation or metabolism, and cell-extrinsic phenomena, such as the host immune response and angiogenesis, have important practical implications for the development of therapeutics as well as guiding the selection of therapies. Agents that target MYC could be highly effective for the treatment of some

cancer types or as individualized treatments for some patients. To date, no direct MYC inhibitors have been approved for clinical use, although we envision that such agents, once identified, will be effective either alone or in combination with other therapeutic agents to treat *MYC*-driven cancers. Furthermore, understanding and targeting those MYC functions that are independent of its transcriptional activity might provide alternative therapeutic strategies. Hence, molecular understanding of how MYC causes cancer should enable efforts to identify which patients with *MYC*-driven cancers are most likely to have a response to specific therapies such as direct MYC inhibitors, MYC synthetic lethal agents, MYC-guided immunotherapies and/or MYC-guided metabolic treatments (FIG. 6). Thus, we envision that patients with tumours could be assigned to MYC^{high} or MYC^{low} phenotypes defined by genomic events including *MYC* amplifications and translocations and/or expression of an MYC activation gene signature^{226,227}. Subsequently, deep phenotyping of tumours using different approaches (including DNA and RNA sequencing, mass cytometry, or mass spectrometry) would guide the selection of therapeutic strategies to target MYC (FIG. 6). For example, patients with tumours harbouring *MYC* amplifications can potentially be treated with direct MYC inhibitors while those with enrichment of synthetic lethal targets of *MYC* would receive drugs targeting these specific gene products. MYC-directed therapies could be used to sensitize MYC^{high} immunologically ‘cold’ tumours to immunotherapies, whereas tumours in which MYC predominantly induces metabolic dysregulation can potentially be treated with drugs targeting metabolic pathways (such as lipid metabolism). In summary, discoveries from the past few years elucidating the specific mechanisms by which MYC drives cancer can potentially enable researchers to develop novel therapeutic strategies to target MYC.

Conclusions

MYC and other genes in the *MYC* superfamily are among the most commonly activated signalling mechanisms in human cancer. MYC initiates and maintains cancer through both tumour cell-intrinsic mechanisms and host immune and TME-dependent mechanisms. In experimental models, MYC inactivation can induce rapid tumour regression through effects on tumour cells, which occur as a consequence of oncogene addiction, as well as the restoration of immune responses. To date, however, the MYC pathway remains to be successfully targeted therapeutically. In this Review, we have described multiple approaches for targeting MYC or MYC-regulated pathways that are being evaluated in clinical trials (Supplementary Table 1). Furthermore, we propose that MYC and other members of the MYC pathway could serve as biomarkers to stratify patients for specific therapies (FIG. 6).

MYC is a particularly exciting therapeutic target because it regulates not only tumour cell-intrinsic growth but also the host immune response. Thus, immune surveillance might be restored by targeting the MYC pathway. For more than 30 years, MYC was presumed to contribute to tumorigenesis through its direct




Fig. 6 | Proposed biomarker-stratified therapeutic strategies to target MYC-driven cancers. Patients with cancer can be assigned to MYC^{hi} and MYC^{low} subgroups defined by enrichment of an MYC activation signature in tumour samples. Patients with MYC-driven tumours can potentially be further classified according to various disease phenotypes on the basis of the dominant mechanism of action of MYC, which would enable stratification to receive different treatments. For example, patients harbouring tumours with MYC amplifications could be treated with direct MYC inhibitors, whereas those with enrichment of synthetic lethal targets of MYC would receive agents targeting these specific gene products. By contrast, MYC-directed therapies could be used to sensitize MYC^{hi} immunologically ‘cold’ tumours to immunotherapies, whereas patients with tumours in which MYC prominently induces metabolic dysregulation can be treated with drugs targeting these pathways. In summary, discoveries elucidating the specific mechanisms by which MYC drives cancer are enabling the development of novel and selective therapeutic strategies to target MYC in human cancers.

effects on cancer cells. Nowadays, the role of MYC as a grand orchestrator of biological processes, not only inherent to tumour cells but also relating to the TME and the host immune system, has become apparent. MYC coordinates intercellular communications that occur between the cancer cells and specific immune cells, thus enabling tumour initiation, progression and metastasis.

Experimentally, targeting MYC and the MYC pathway seems to both affect cancer cells and restore antitumour immunity. The evidence of MYC as a grand orchestrator provides further impetus to develop therapeutics to target this otherwise elusive oncogenic pathway.

Published online 10 September 2021

- Meyer, N. & Penn, L. Z. Reflecting on 25 years with MYC. *Nat. Rev. Cancer* **8**, 976–990 (2008).
- Dang, C. V. MYC on the path to cancer. *Cell* **149**, 22–35 (2012).
- Schaub, F. X. et al. Pan-cancer alterations of the MYC oncogene and its proximal network across The Cancer Genome Atlas. *Cell Syst.* **6**, 282–300.e2 (2018).
- Kalkat, M. et al. MYC deregulation in primary human cancers. *Genes* **8**, 151 (2017).
- Kress, T. R., Sabò, A. & Amati, B. MYC: connecting selective transcriptional control to global RNA production. *Nat. Rev. Cancer* **15**, 593–607 (2015).
- Conacci-Sorrell, M., McFerrin, L. & Eisenman, R. N. An overview of MYC and its interactome. *Cold Spring Harb. Perspect. Med.* **4**, a014357 (2014).
- Williamson, D. et al. Relationship between MYCN copy number and expression in rhabdomyosarcomas and correlation with adverse prognosis in the alveolar subtype. *J. Clin. Oncol.* **23**, 880–888 (2005).
- Williams, R. D. et al. Molecular profiling reveals frequent gain of MYCN and anaplasia-specific loss of 4q and 14q in Wilms tumor. *Genes Chromosomes Cancer* **50**, 982–995 (2011).
- Berger, A. et al. N-Myc-mediated epigenetic reprogramming drives lineage plasticity in advanced

- prostate cancer. *J. Clin. Invest.* **129**, 3924–3940 (2019).
10. Rickman, D. S., Schulte, J. H. & Eilers, M. The expanding world of N-MYC–driven tumors. *Cancer Discov.* **8**, 150–163 (2018).
 11. Cheng, J. et al. Merkel cell polyomavirus recruits MYCL to the EP400 complex to promote oncogenesis. *PLoS Pathog.* **13**, e1006668 (2017).
 12. Ohshima, K. et al. Integrated analysis of gene expression and copy number identified potential cancer driver genes with amplification-dependent overexpression in 1,454 solid tumors. *Sci. Rep.* **7**, 641 (2017).
 13. Boutros, P. C. et al. Spatial genomic heterogeneity within localized, multifocal prostate cancer. *Nat. Genet.* **47**, 736–745 (2015).
 14. Adams, J. M. et al. The c-myc oncogene driven by immunoglobulin enhancers induces lymphoid malignancy in transgenic mice. *Nature* **318**, 533–538 (1985).
 15. Dalla-Favera, R. et al. Human c-myc onc gene is located on the region of chromosome 8 that is translocated in Burkitt lymphoma cells. *Proc. Natl Acad. Sci. USA* **79**, 7824–7827 (1982).
 16. Gurel, B. et al. Nuclear MYC protein overexpression is an early alteration in human prostate carcinogenesis. *Mod. Pathol.* **21**, 1156–1167 (2008).
 17. Stock, C., Kager, L., Fink, F. M., Gadner, H. & Ambros, P. F. Chromosomal regions involved in the pathogenesis of osteosarcomas. *Genes Chromosomes Cancer* **28**, 329–336 (2000).
 18. Jain, M. et al. Sustained loss of a neoplastic phenotype by brief inactivation of MYC. *Science* **297**, 102–104 (2002).
 19. Shroff, E. H. et al. MYC oncogene overexpression drives renal cell carcinoma in a mouse model through glutamine metabolism. *Proc. Natl Acad. Sci. USA* **112**, 6539–6544 (2015).
 20. Felsner, D. W. & Bishop, J. M. Reversible tumorigenesis by MYC in hematopoietic lineages. *Mol. Cell* **4**, 199–207 (1999).
 21. Wu, C.-H. et al. Cellular senescence is an important mechanism of tumor regression upon c-Myc inactivation. *Proc. Natl Acad. Sci. USA* **104**, 13028–13033 (2007).
 22. Lourenco, C. et al. Modelling the MYC-driven normal-to-tumour switch in breast cancer. *Dis. Model. Mech.* **12**, dmm038083 (2019).
 23. Sodir, N. M. et al. MYC instructs and maintains pancreatic adenocarcinoma phenotype. *Cancer Discov.* **10**, 588–607 (2020).
 24. Soucek, L. et al. Inhibition of Myc family proteins eradicates KRas-driven lung cancer in mice. *Genes Dev.* **27**, 504–513 (2013).
 25. Jung, L. A. et al. OmoMYC blunts promoter invasion by oncogenic MYC to inhibit gene expression characteristic of MYC-dependent tumors. *Oncogene* **36**, 1911–1924 (2017).
 26. Demma, M. J. et al. Omomyc reveals new mechanisms to inhibit the MYC oncogene. *Mol. Cell. Biol.* **39**, e00248-19 (2019).
 27. Weinstein, I. B. Cancer: addiction to oncogenes — the Achilles heal of cancer. *Science* **297**, 63–64 (2002).
 28. Hanahan, D. & Weinberg, R. A. Hallmarks of cancer: the next generation. *Cell* **144**, 646–674 (2011).
 29. Carroll, P. A., Freie, B. W., Mathysaraja, H. & Eisenman, R. N. The MYC transcription factor network: balancing metabolism, proliferation and oncogenesis. *Front. Med.* **12**, 412–425 (2018).
 30. Dang, C. V. c-Myc target genes involved in cell growth, apoptosis, and metabolism. *Mol. Cell. Biol.* **19**, 1–11 (1999).
 31. Casey, S. C., Baylot, V. & Felsner, D. W. The MYC oncogene is a global regulator of the immune response. *Blood* **131**, 2007–2015 (2018).
 32. Casey, S. C., Baylot, V. & Felsner, D. W. MYC: master regulator of immune privilege. *Trends Immunol.* **38**, 298–305 (2017).
 33. Casacuberta-Serra, S. & Soucek, L. Myc and Ras, the Bonnie and Clyde of immune evasion. *Transl. Cancer Res.* **7** (Suppl. 4), S457–S459 (2018).
 34. Baluapuri, A., Wolf, E. & Eilers, M. Target gene-independent functions of MYC oncoproteins. *Nat. Rev. Mol. Cell Biol.* **21**, 255–267 (2020).
 35. Stine, Z. E., Walton, Z. E., Altman, B. J., Hsieh, A. L. & Dang, C. V. MYC, metabolism, and cancer. *Cancer Discov.* **5**, 1024–1039 (2015).
 36. Bernard, S. & Eilers, M. Control of cell proliferation and growth by Myc proteins. *Results Probl. Cell. Differ.* **42**, 329–342 (2006).
 37. Bretones, G., Delgado, M. D. & León, J. Myc and cell cycle control. *Biochim. Biophys. Acta* **1849**, 506–516 (2015).
 38. Levens, D. 'You don't muck with MYC'. *Genes Cancer* **1**, 547–554 (2010).
 39. Beroukhi, R. et al. The landscape of somatic copy-number alteration across human cancers. *Nature* **463**, 899–905 (2010).
 40. Turner, K. M. et al. Extrachromosomal oncogene amplification drives tumour evolution and genetic heterogeneity. *Nature* **543**, 122–125 (2017).
 41. Wu, S. et al. Circular ecDNA promotes accessible chromatin and high oncogene expression. *Nature* **575**, 699–703 (2019).
 42. Boxer, L. M. & Dang, C. V. Translocations involving c-myc and c-myc function. *Oncogene* **20**, 5595–5610 (2001).
 43. Drach, J. et al. The biology of multiple myeloma. *J. Cancer Res. Clin. Oncol.* **126**, 441–447 (2000).
 44. Dudley, J. P., Mertz, J. A., Rajan, L., Lozano, M. & Broussard, D. R. What retroviruses teach us about the involvement of c-Myc in leukemias and lymphomas. *Leukemia* **16**, 1086–1098 (2002).
 45. Weng, A. P. et al. c-Myc is an important direct target of Notch1 in T-cell acute lymphoblastic leukemia/lymphoma. *Genes Dev.* **20**, 2096–2109 (2006).
 46. He, T. C. et al. Identification of c-MYC as a target of the APC pathway. *Science* **281**, 1509–1512 (1998).
 47. Yagi, K. et al. c-myc is a downstream target of the Smad pathway. *J. Biol. Chem.* **277**, 854–861 (2002).
 48. Allen-Petersen, B. L. & Sears, R. C. Mission possible: advances in MYC therapeutic targeting in cancer. *BioDrugs* **33**, 539–553 (2019).
 49. Sears, R. et al. Multiple Ras-dependent phosphorylation pathways regulate Myc protein stability. *Genes Dev.* **14**, 2501–2514 (2000).
 50. Arnold, H. K. & Sears, R. C. Protein phosphatase 2A regulatory subunit B56alpha associates with c-myc and negatively regulates c-myc accumulation. *Mol. Cell. Biol.* **26**, 2832–2844 (2006).
 51. Farrell, A. S. et al. MYC regulates ductal-neuroendocrine lineage plasticity in pancreatic ductal adenocarcinoma associated with poor outcome and chemoresistance. *Nat. Commun.* **8**, 1728 (2017).
 52. Wang, X. et al. Phosphorylation regulates c-Myc's oncogenic activity in the mammary gland. *Cancer Res.* **71**, 925–936 (2011).
 53. Reavie, L. et al. Regulation of c-Myc ubiquitination controls chronic myelogenous leukemia initiation and progression. *Cancer Cell* **23**, 362–375 (2013).
 54. Ruvoilo, P. P. The broken 'Off' switch in cancer signaling: PP2A as a regulator of tumorigenesis, drug resistance, and immune surveillance. *BBA Clin.* **6**, 87–99 (2016).
 55. Kauko, O. et al. PP2A inhibition is a druggable MEK inhibitor resistance mechanism in KRAS-mutant lung cancer cells. *Sci. Transl. Med.* **10**, eaaq1093 (2018).
 56. Zhou, X. Z. & Lu, K. P. The isomerase PIN1 controls numerous cancer-driving pathways and is a unique drug target. *Nat. Rev. Cancer* **16**, 463–478 (2016).
 57. Cheng, C.-W. & Tse, E. PIN1 in cell cycle control and cancer. *Front. Pharmacol.* **9**, 1367 (2018).
 58. King, B. et al. The ubiquitin ligase FBXW7 modulates leukemia-initiating cell activity by regulating MYC stability. *Cell* **153**, 1552–1566 (2013).
 59. Yeh, C.-H., Bellon, M. & Nicot, C. FBXW7: a critical tumor suppressor of human cancers. *Mol. Cancer* **17**, 115 (2018).
 60. Cao, J., Ge, M.-H. & Ling, Z.-Q. FBXw7 tumor suppressor: a vital regulator contributes to human tumorigenesis. *Medicine* **95**, e2496 (2016).
 61. Grandori, C., Cowley, S. M., James, L. P. & Eisenman, R. N. The Myc/Max/Mad network and the transcriptional control of cell behavior. *Annu. Rev. Cell Dev. Biol.* **16**, 653–699 (2000).
 62. Edelmann, J. et al. Frequent evolution of copy number alterations in CLL following first-line treatment with FC(R) is enriched with TP53 alterations: results from the CLL8 trial. *Leukemia* **31**, 734–738 (2017).
 63. Yang, G. & Hurlin, P. J. MNT and emerging concepts of MNT-MYC antagonism. *Genes* **8**, 83 (2017).
 64. Fredlund, E., Ringnér, M., Maris, J. M. & Pålman, S. High MYC pathway activity and low stage of neuronal differentiation associate with poor outcome in neuroblastoma. *Proc. Natl Acad. Sci. USA* **105**, 14094–14099 (2008).
 65. Diolaiti, D., McFerrin, L., Carroll, P. A. & Eisenman, R. N. Functional interactions among members of the MAX and MLX transcriptional network during oncogenesis. *Biochim. Biophys. Acta* **1849**, 484–500 (2015).
 66. Gabay, M., Li, Y. & Felsner, D. W. MYC activation is a hallmark of cancer initiation and maintenance. *Cold Spring Harb. Perspect. Med.* **4**, a014241 (2014).
 67. Bailey, S. T. et al. MYC activation cooperates with Vhl and Ink4a/Arf loss to induce clear cell renal cell carcinoma. *Nat. Commun.* **8**, 15770 (2017).
 68. Vaux, D. L., Cory, S. & Adams, J. M. Bcl-2 gene promotes haemopoietic cell survival and cooperates with c-myc to immortalize pre-B cells. *Nature* **335**, 440–442 (1988).
 69. Teitz, T. et al. Caspase 8 is deleted or silenced preferentially in childhood neuroblastomas with amplification of MYCN. *Nat. Med.* **6**, 529–535 (2000).
 70. Wu, K. J. et al. Direct activation of TERT transcription by c-MYC. *Nat. Genet.* **21**, 220–224 (1999).
 71. Hofmann, J. W. et al. Reduced expression of MYC increases longevity and enhances healthspan. *Cell* **160**, 477–488 (2015).
 72. Seton-Rogers, S. Driving immune evasion. *Nat. Rev. Cancer* **18**, 67–67 (2018).
 73. Chappell, J. & Dalton, S. Roles for MYC in the establishment and maintenance of pluripotency. *Cold Spring Harb. Perspect. Med.* **3**, a014381 (2013).
 74. Pelengaris, S. & Khan, M. The many faces of c-MYC. *Arch. Biochem. Biophys.* **416**, 129–136 (2003).
 75. de la Cova, C., Abril, M., Bellosta, P., Gallant, P. & Johnston, L. A. Drosophila myc regulates organ size by inducing cell competition. *Cell* **117**, 107–116 (2004).
 76. Bettess, M. D. et al. c-Myc is required for the formation of intestinal crypts but dispensable for homeostasis of the adult intestinal epithelium. *Mol. Cell. Biol.* **25**, 7868–7878 (2005).
 77. Evan, G. I. et al. Induction of apoptosis in fibroblasts by c-myc protein. *Cell* **69**, 119–128 (1992).
 78. Shi, Y. et al. Role for c-myc in activation-induced apoptotic cell death in T cell hybridomas. *Science* **257**, 212–214 (1992).
 79. Kaczmarek, L., Hyland, J. K., Watt, R., Rosenberg, M. & Baserga, R. Microinjected c-myc as a competence factor. *Science* **228**, 1313–1315 (1985).
 80. Zindy, F. et al. Myc signaling via the ARF tumor suppressor regulates p53-dependent apoptosis and immortalization. *Genes Dev.* **12**, 2424–2433 (1998).
 81. Karn, J., Watson, J. V., Lowe, A. D., Green, S. M. & Vedeckis, W. Regulation of cell cycle duration by c-myc levels. *Oncogene* **4**, 773–787 (1989).
 82. Rosenwald, I. B. Upregulated expression of the genes encoding translation initiation factors eIF-4E and eIF-2a in transformed cells. *Cancer Lett.* **102**, 113–123 (1996).
 83. Boon, K. et al. N-myc enhances the expression of a large set of genes functioning in ribosome biogenesis and protein synthesis. *EMBO J.* **20**, 1383–1393 (2001).
 84. Schmidt, E. V. The role of c-myc in cellular growth control. *Oncogene* **18**, 2988–2996 (1999).
 85. Grewal, S. S., Li, L., Orian, A., Eisenman, R. N. & Edgar, B. A. Myc-dependent regulation of ribosomal RNA synthesis during Drosophila development. *Nat. Cell Biol.* **7**, 295–302 (2005).
 86. Dong, Y., Tu, R., Liu, H. & Qing, G. Regulation of cancer cell metabolism: oncogenic MYC in the driver's seat. *Signal Transduct. Target. Ther.* **5**, 124 (2020).
 87. Kim, J.-W. et al. Evaluation of myc E-box phylogenetic footprints in glycolytic genes by chromatin immunoprecipitation assays. *Mol. Cell. Biol.* **24**, 5923–5936 (2004).
 88. Osthur, R. C. et al. Deregulation of glucose transporter 1 and glycolytic gene expression by c-Myc. *J. Biol. Chem.* **275**, 21797–21800 (2000).
 89. Wise, D. R. et al. Myc regulates a transcriptional program that stimulates mitochondrial glutaminolysis and leads to glutamine addiction. *Proc. Natl Acad. Sci. USA* **105**, 18782–18787 (2008).
 90. Dominguez-Sola, D. et al. Non-transcriptional control of DNA replication by c-Myc. *Nature* **448**, 445–451 (2007).
 91. Takahashi, K. & Yamanaka, S. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* **126**, 663–676 (2006).
 92. García-Gutiérrez, L., Delgado, M. D. & León, J. MYC oncogene contributions to release of cell cycle brakes. *Genes* **10**, 244 (2019).
 93. Li, Z. et al. c-Myc suppression of DNA double-strand break repair. *Neoplasia* **14**, 1190–1202 (2012).
 94. Felsner, D. W. & Bishop, J. M. Transient excess of MYC activity can elicit genomic instability and tumorigenesis. *Proc. Natl Acad. Sci. USA* **96**, 3940–3944 (1999).

95. Kuzyk, A. & Mai, S. c-MYC-induced genomic instability. *Cold Spring Harb. Perspect. Med.* **4**, a014373 (2014).
96. Karlsson, A. et al. Defective double-strand DNA break repair and chromosomal translocations by MYC overexpression. *Proc. Natl Acad. Sci. USA* **100**, 9974–9979 (2003).
97. Vafa, O. et al. c-Myc can induce DNA damage, increase reactive oxygen species, and mitigate p53 function: a mechanism for oncogene-induced genetic instability. *Mol. Cell* **9**, 1031–1044 (2002).
98. Dang, C. V., Li, F. & Lee, L. A. Could MYC induction of mitochondrial biogenesis be linked to ROS production and genomic instability? *Cell Cycle* **4**, 1465–1466 (2005).
99. Reimann, M. et al. The Myc-evoked DNA damage response accounts for treatment resistance in primary lymphomas in vivo. *Blood* **110**, 2996–3004 (2007).
100. Beer, S. et al. Developmental context determines latency of MYC-induced tumorigenesis. *PLoS Biol.* **2**, e352 (2004).
101. Ray, S. et al. MYC can induce DNA breaks in vivo and in vitro independent of reactive oxygen species. *Cancer Res.* **66**, 6598–6605 (2006).
102. Baudino, T. A. et al. c-Myc is essential for vasculogenesis and angiogenesis during development and tumor progression. *Genes Dev.* **16**, 2530–2543 (2002).
103. Shchors, K. et al. The Myc-dependent angiogenic switch in tumors is mediated by interleukin 1 β . *Genes Dev.* **20**, 2527–2538 (2006).
104. Dews, M. et al. The Myc–miR-17-92 axis blunts TGF β signaling and production of multiple TGF β -dependent antiangiogenic factors. *Cancer Res.* **70**, 8233–8246 (2010).
105. Giuriato, S. et al. Sustained regression of tumors upon MYC inactivation requires p53 or thrombospondin-1 to reverse the angiogenic switch. *Proc. Natl Acad. Sci. USA* **103**, 16266–16271 (2006).
106. Rakhra, K. et al. CD4⁺ T cells contribute to the remodeling of the microenvironment required for sustained tumor regression upon oncogene inactivation. *Cancer Cell* **18**, 485–498 (2010).
107. Kortlever, R. M. et al. Myc cooperates with Ras by programming inflammation and immune suppression. *Cell* **171**, 1301–1315.e14 (2017).
108. Casey, S. C. et al. MYC regulates the antitumor immune response through CD47 and PD-L1. *Science* **352**, 227–231 (2016).
109. Swaminathan, S. et al. MYC functions as a switch for natural killer cell-mediated immune surveillance of lymphoid malignancies. *Nat. Commun.* **11**, 2860 (2020).
110. Maeda, T. et al. MUC1-C induces PD-L1 and immune evasion in triple-negative breast cancer. *Cancer Res.* **78**, 205–215 (2018).
111. Muthalagu, N. et al. Repression of the type I interferon pathway underlies MYC & KRAS-dependent evasion of NK & B cells in pancreatic ductal adenocarcinoma. *Cancer Discov.* **10**, 872–887 (2020).
112. Dhanasekaran, R. et al. MYC and Twist1 cooperate to drive metastasis by eliciting crosstalk between cancer and innate immunity. *eLife* **9**, e50731 (2020).
113. Bernards, R., Dessain, S. K. & Weinberg, R. A. N-myc amplification causes down-modulation of MHC class I antigen expression in neuroblastoma. *Cell* **47**, 667–674 (1986).
114. Braun, J., Felsner, D. W. & Goodglick, L. A. c-myc, MHCI, and NK resistance in immunodeficiency lymphomas. *Ann. N. Y. Acad. Sci.* **651**, 467–469 (1992).
115. Versteeg, R., Noordermeer, I. A., Krüse-Wolters, M., Ruiters, D. J. & Schrier, P. I. c-myc down-regulates class I HLA expression in human melanomas. *EMBO J.* **7**, 1023–1029 (1988).
116. van Riggelen, J. et al. The interaction between Myc and Miz1 is required to antagonize TGF β -dependent autocrine signaling during lymphoma formation and maintenance. *Genes Dev.* **24**, 1281–1294 (2010).
117. Reimann, M. et al. Tumor stroma-derived TGF- β limits myc-driven lymphomagenesis via Suv39h1-dependent senescence. *Cancer Cell* **17**, 262–272 (2010).
118. Xu, Y. et al. Translation control of the immune checkpoint in cancer and its therapeutic targeting. *Nat. Med.* **25**, 301–311 (2019).
119. Vartuli, R. L. et al. Eya3 promotes breast tumor-associated immune suppression via threonine phosphatase-mediated PD-L1 upregulation. *J. Clin. Invest.* **128**, 2535–2550 (2018).
120. Dhanasekaran, R. et al. MYC and Twist1 cooperate to drive metastasis by eliciting crosstalk between cancer and innate immunity. *eLife* **9**, e50731 (2020).
121. Shachaf, C. M. et al. MYC inactivation uncovers pluripotent differentiation and tumour dormancy in hepatocellular cancer. *Nature* **431**, 1112–1117 (2004).
122. Tran, P. T. et al. Combined inactivation of MYC and K-Ras oncogenes reverses tumorigenesis in lung adenocarcinomas and lymphomas. *PLoS One* **3**, e2125 (2008).
123. Tran, P. T. et al. Survival and death signals can predict tumor response to therapy after oncogene inactivation. *Sci. Transl. Med.* **3**, 103ra99 (2011).
124. Bui, T. V. & Mendell, J. T. Myc: maestro of microRNAs. *Genes Cancer* **1**, 568–575 (2010).
125. Li, Y., Choi, P. S., Casey, S. C., Dill, D. L. & Felsner, D. W. MYC through miR-17-92 Suppresses specific target genes to maintain survival, autonomous proliferation, and a neoplastic state. *Cancer Cell* **26**, 262–272 (2014).
126. Chang, T.-C. et al. Widespread microRNA repression by Myc contributes to tumorigenesis. *Nat. Genet.* **40**, 43–50 (2008).
127. Adams, C. M., Hiebert, S. W. & Eisen, C. M. Myc induces miRNA-mediated apoptosis in response to HDAC inhibition in hematologic malignancies. *Cancer Res.* **76**, 736–748 (2016).
128. Zhou, L. et al. Silencing of thrombospondin-1 is critical for myc-induced metastatic phenotypes in medulloblastoma. *Cancer Res.* **70**, 8199–8210 (2010).
129. Swaminathan, S. et al. MYC functions as a master switch for natural killer cell-mediated immune surveillance of lymphoid malignancies. *Blood* **132**, 2619–2619 (2018).
130. Casey, S. C., Li, Y., Fan, A. C. & Felsner, D. W. Oncogene withdrawal engages the immune system to induce sustained cancer regression. *J. Immunother. Cancer* **2**, 24 (2014).
131. Sharma, S. V., Fischbach, M. A., Haber, D. A. & Settleman, J. ‘Oncogenic shock’: explaining oncogene addiction through differential signal attenuation. *Clin. Cancer Res.* **12**, 4392s–4395s (2006).
132. Fisher, G. H. et al. Induction and apoptotic regression of lung adenocarcinomas by regulation of a K-Ras transgene in the presence and absence of tumor suppressor genes. *Genes Dev.* **15**, 3249–3262 (2001).
133. Kohl, N. E. et al. Protein farnesyltransferase inhibitors block the growth of ras-dependent tumors in nude mice. *Proc. Natl Acad. Sci. USA* **91**, 9141–9145 (1994).
134. Moody, S. E. et al. Conditional activation of Neu in the mammary epithelium of transgenic mice results in reversible pulmonary metastasis. *Cancer Cell* **2**, 451–461 (2002).
135. Hori, S. S. et al. A mathematical model of tumor regression and recurrence after therapeutic oncogene inactivation. *Sci. Rep.* **11**, 1341 (2021).
136. Soucek, L. et al. Mast cells are required for angiogenesis and macroscopic expansion of Myc-induced pancreatic islet tumors. *Nat. Med.* **13**, 1211–1218 (2007).
137. Mahaud-Fernandez, W. D. & Felsner, D. W. The Myc and Ras partnership in cancer: indistinguishable alliance or contextual relationship? *Cancer Res.* **80**, 3799–3802 (2020).
138. Sun, L. et al. Gastric cancer mesenchymal stem cells derived IL-8 induces PD-L1 expression in gastric cancer cells via STAT3/mTOR-c-Myc signal axis. *Cell Death Dis.* **9**, 928 (2018).
139. Kharma, B. et al. STAT1 drives tumor progression in serous papillary endometrial cancer. *Cancer Res.* **74**, 6519–6530 (2014).
140. Topper, M. J. et al. Epigenetic therapy Ties MYC depletion to reversing immune evasion and treating lung cancer. *Cell* **171**, 1284–1300.e21 (2017).
141. Coelho, M. A. et al. Oncogenic RAS signaling promotes tumor immunoresistance by stabilizing PD-L1 mRNA. *Immunity* **47**, 1083–1099.e6 (2017).
142. Melaiu, O. et al. PD-L1 is a therapeutic target of the bromodomain inhibitor JQ1 and, combined with HLA Class I, a promising prognostic biomarker in neuroblastoma. *Clin. Cancer Res.* **23**, 4462–4472 (2017).
143. Atsaves, V. et al. PD-L1 is commonly expressed and transcriptionally regulated by STAT3 and MYC in ALK-negative anaplastic large-cell lymphoma. *Leukemia* **31**, 1633–1637 (2017).
144. Pai, S. et al. CD47-SIRP α signaling induces epithelial-mesenchymal transition and cancer stemness and links to a poor prognosis in patients with oral squamous cell carcinoma. *Cells* **8**, 1658 (2019).
145. Kaur, S. et al. Thrombospondin-1 signaling through CD47 inhibits self-renewal by regulating c-Myc and other stem cell transcription factors. *Sci. Rep.* **3**, 1673 (2013).
146. Li, W. et al. Targeting MYC activity in double-hit lymphoma with MYC and BCL2 and/or BCL6 rearrangements with epigenetic bromodomain inhibitors. *J. Hematol. Oncol.* **12**, 73 (2019).
147. Kamijo, H. et al. Thrombospondin-1 promotes tumor progression in cutaneous T-cell lymphoma via CD47. *Leukemia* **34**, 845–856 (2020).
148. Felsner, D. W., Rhim, S. H. & Braun, J. A murine model for B-cell lymphomagenesis in immunocompromised hosts: natural killer cells are an important component of host resistance to premalignant B-cell lines. *Cancer Res.* **50**, 7050–7056 (1990).
149. El-Jawhari, J. J. et al. Blocking oncogenic RAS enhances tumor cell surface MHC class I expression but does not alter susceptibility to cytotoxic lymphocytes. *Mol. Immunol.* **58**, 160–168 (2014).
150. Ji, H. et al. K-ras activation generates an inflammatory response in lung tumors. *Oncogene* **25**, 2105–2112 (2006).
151. Van Dang, C. & Kim, J.-W. Convergence of cancer metabolism and immunity: an overview. *Biomol. Ther.* **26**, 4–9 (2018).
152. Gouw, A. M., Hsieh, A. L., Stine, Z. E. & Dang, C. V. In: *Tumor Cell Metabolism* (Mazurek S., Shoshan M. eds) 101–122 (Springer, 2015).
153. Dang, C. V. MYC, metabolism, cell growth, and tumorigenesis. *Cold Spring Harb. Perspect. Med.* **3**, a014217 (2013).
154. Gouw, A. M. et al. The MYC oncogene cooperates with sterol-regulated element-binding protein to regulate lipogenesis essential for neoplastic growth. *Cell Metab.* **30**, 556–572.e5 (2019).
155. Ricciardi, S. et al. The translational machinery of human CD4 T cells is poised for activation and controls the switch from quiescence to metabolic remodeling. *Cell Metab.* **28**, 895–906.e5 (2018).
156. Friesen, L. R., Gu, B. & Kim, C. H. A ligand-independent fast function of RAR α promotes exit from metabolic quiescence upon T cell activation and controls T cell differentiation. *Mucosal Immunol.* **14**, 100–112 (2021).
157. Chapman, N. M., Boothby, M. R. & Chi, H. Metabolic coordination of T cell quiescence and activation. *Nat. Rev. Immunol.* **20**, 55–70 (2020).
158. Kapur, S. & Pal, A. Immune cell activation: stimulation, costimulation, and regulation of cellular activation. *Immune Response Activation Immunodiol.* <https://doi.org/10.5772/intechopen.81568> (2019).
159. Shim, H. et al. c-Myc transactivation of LDH-A: implications for tumor metabolism and growth. *Proc. Natl Acad. Sci. USA* **94**, 6658–6663 (1997).
160. Briggs, K. J. et al. Paracrine Induction of HIF by glutamate in breast cancer: EglN1 senses cysteine. *Cell* **166**, 126–139 (2016).
161. Bott, A. J. et al. Oncogenic Myc induces expression of glutamine synthetase through promoter demethylation. *Cell Metab.* **22**, 1068–1077 (2015).
162. Gerriets, V. A. & Rathmell, J. C. Metabolic pathways in T cell fate and function. *Trends Immunol.* **33**, 168–173 (2012).
163. Verbist, K. C. et al. Metabolic maintenance of cell asymmetry following division in activated T lymphocytes. *Nature* **532**, 389–393 (2016).
164. Le, A. et al. Glucose-independent glutamine metabolism via TCA cycling for proliferation and survival in B cells. *Cell Metab.* **15**, 110–121 (2012).
165. Eberlin, L. S. et al. Alteration of the lipid profile in lymphomas induced by MYC overexpression. *Proc. Natl Acad. Sci. USA* **111**, 10450–10455 (2014).
166. Allison, K. E., Coomber, B. L. & Bridle, B. W. Metabolic reprogramming in the tumour microenvironment: a hallmark shared by cancer cells and T lymphocytes. *Immunology* **152**, 175–184 (2017).
167. Koudhi, S., Ben Ayed, F. & Elgaaiel, A. B. Targeting tumor metabolism: a new challenge to improve immunotherapy. *Front. Immunol.* **9**, 353 (2018).
168. Massó-Vallés, D. & Soucek, L. Blocking myc to treat cancer: reflecting on two decades of omomyc. *Cells* **9**, 883 (2020).
169. Wolf, E. & Eilers, M. Targeting MYC proteins for tumor therapy. *Annu. Rev. Cancer Biol.* **4**, 61–75 (2020).
170. McKeown, M. R. & Bradner, J. E. Therapeutic strategies to inhibit MYC. *Cold Spring Harb. Perspect. Med.* **4**, a014266 (2014).
171. Fletcher, S. & Prochownik, E. V. Small-molecule inhibitors of the Myc oncoprotein. *Biochim. Biophys. Acta* **1849**, 525–543 (2015).

172. Whitfield, J. R., Beaulieu, M.-E. & Soucek, L. Strategies to inhibit myc and their clinical applicability. *Front. Cell Dev. Biol.* **5**, 10 (2017).
173. Carabet, L. A., Rennie, P. S. & Cherkasov, A. Therapeutic inhibition of myc in cancer: structural bases and computer-aided drug discovery approaches. *Int. J. Mol. Sci.* **20**, 120 (2018).
174. Madden, S. K., de Araujo, A. D., Gerhardt, M., Fairlie, D. P. & Mason, J. M. Taking the Myc out of cancer: toward therapeutic strategies to directly inhibit c-Myc. *Mol. Cancer* **20**, 3 (2021).
175. Thng, D. K. H., Toh, T. B. & Chow, E. K.-H. Capitalizing on synthetic lethality of MYC to treat cancer in the digital age. *Trends Pharmacol. Sci.* **42**, 166–182 (2021).
176. Cascon, A. & Robledo, M. MAX and MYC: a heritable breakup. *Cancer Res.* **72**, 3119–3124 (2012).
177. Han, H. et al. Small-molecule MYC inhibitors suppress tumor growth and enhance immunotherapy. *Cancer Cell* **36**, 483–497.e15 (2019).
178. Struntz, N. B. et al. Stabilization of the max homodimer with a small molecule attenuates Myc-driven transcription. *Cell Chem. Biol.* **26**, 711–725.e14 (2019).
179. Lustig, L. C. et al. Inhibiting MYC binding to the E-box DNA motif by ME47 decreases tumour xenograft growth. *Oncogene* **36**, 6830–6837 (2017).
180. Boike, L. et al. Discovery of a functional covalent ligand targeting an intrinsically disordered cysteine within MYC. *Cell Chem. Biol.* **28**, 4–13.e17 (2021).
181. Pourdehnad, M. et al. Myc and mTOR converge on a common node in protein synthesis control that confers synthetic lethality in Myc-driven cancers. *Proc. Natl Acad. Sci. USA* **110**, 11988–11993 (2013).
182. Wiegner, A. et al. Targeting translation initiation bypasses signaling crosstalk mechanisms that maintain high MYC levels in colorectal cancer. *Cancer Discov.* **5**, 768–781 (2015).
183. Gustafson, W. C. et al. Drugging MYCN through an allosteric transition in Aurora kinase A. *Cancer Cell* **26**, 414–427 (2014).
184. Yang, D. et al. Therapeutic potential of a synthetic lethal interaction between the MYC proto-oncogene and inhibition of aurora-B kinase. *Proc. Natl Acad. Sci. USA* **107**, 13836–13841 (2010).
185. Xiao, D. et al. Polo-like kinase-1 regulates myc stabilization and activates a feedforward circuit promoting tumor cell survival. *Mol. Cell* **64**, 493–506 (2016).
186. Yeh, E. et al. A signalling pathway controlling c-Myc degradation that impacts oncogenic transformation of human cells. *Nat. Cell Biol.* **6**, 308–318 (2004).
187. Farrell, A. S. et al. Pin1 regulates the dynamics of c-Myc DNA binding to facilitate target gene regulation and oncogenesis. *Mol. Cell. Biol.* **33**, 2930–2949 (2013).
188. Winter, G. E. et al. Phthalimide conjugation as a strategy for in vivo target protein degradation. *Science* **348**, 1376–1381 (2015).
189. Bondeson, D. P. et al. Catalytic in vivo protein knockdown by small-molecule PROTACs. *Nat. Chem. Biol.* **11**, 611–617 (2015).
190. Koh, C. M., Sabo, A. & Guccione, E. Targeting MYC in cancer therapy: RNA processing offers new opportunities. *Bioessays* **38**, 266–275 (2016).
191. Ricker, J. L., Mata, J. E., Iversen, P. L. & Gattone, V. H. c-myc antisense oligonucleotide treatment ameliorates murine ARPKD. *Kidney Int.* **61** (Suppl. 1), S125–S131 (2002).
192. Yu, B. W., Nguyen, D., Anderson, S. & Allegra, C. A. Phosphorothioated antisense c-myc oligonucleotide inhibits the growth of human colon carcinoma cells. *Anticancer Res.* **17**, 4407–4413 (1997).
193. Cotter, F. E. Antisense oligonucleotides: considerations for lymphoma therapy. *Hematology* **1**, 53–57 (1996).
194. Dhanasekaran, R. et al. MYC ASO impedes tumorigenesis and elicits oncogene addiction in autochthonous transgenic mouse models of HCC and RCC. *Mol. Ther. Nucleic Acids* **21**, 850–859 (2020).
195. Filippakopoulos, P. et al. Selective inhibition of BET bromodomains. *Nature* **468**, 1067–1073 (2010).
196. Raina, K. et al. PROTAC-induced BET protein degradation as a therapy for castration-resistant prostate cancer. *Proc. Natl Acad. Sci. USA* **113**, 7124–7129 (2016).
197. Felsenstein, K. M. et al. Small molecule microarrays enable the identification of a selective, quadruplex-binding inhibitor of MYC expression. *ACS Chem. Biol.* **11**, 139–148 (2016).
198. Brown, R. V., Danford, F. L., Gokhale, V., Hurley, L. H. & Brooks, T. A. Demonstration that drug-targeted down-regulation of MYC in non-Hodgkins lymphoma is directly mediated through the promoter G-quadruplex. *J. Biol. Chem.* **286**, 41018–41027 (2011).
199. Dutta, D. et al. Cell penetrating thiazole peptides inhibit c-MYC expression via site-specific targeting of c-MYC G-quadruplex. *Nucleic Acids Res.* **46**, 5355–5365 (2018).
200. Hu, M.-H. et al. Discovery of a new four-leaf clover-like ligand as a potent c-MYC transcription inhibitor specifically targeting the promoter G-quadruplex. *J. Med. Chem.* **61**, 2447–2459 (2018).
201. Panda, D., Saha, P., Das, T. & Dash, J. Target guided synthesis using DNA nano-templates for selectively assembling a G-quadruplex binding c-MYC inhibitor. *Nat. Commun.* **8**, 16103 (2017).
202. Harbor, C. S., Dang, C. V. & Eisenman, R. N. *Myc and the pathway to cancer* (Cold Spring Harbor Laboratory Press, 2014).
203. Harrison, C. Synthetic lethality enables targeting of MYC. *Nat. Rev. Drug Discov.* **11**, 602–602 (2012).
204. Kessler, J. D. et al. A SUMOylation-dependent transcriptional subprogram is required for Myc-driven tumorigenesis. *Science* **335**, 348–353 (2012).
205. Wang, Y. et al. Synthetic lethal targeting of MYC by activation of the DR5 death receptor pathway. *Cancer Cell* **5**, 501–512 (2004).
206. Chen, L. et al. CRISPR-Cas9 screen reveals a MYCN-amplified neuroblastoma dependency on EZH2. *J. Clin. Invest.* **128**, 446–462 (2018).
207. Chipumuro, E. et al. CDK7 inhibition suppresses super-enhancer-linked oncogenic transcription in MYCN-driven cancer. *Cell* **159**, 1126–1139 (2014).
208. Kwiatkowski, N. et al. Targeting transcription regulation in cancer with a covalent CDK7 inhibitor. *Nature* **511**, 616–620 (2014).
209. Krüger, K. et al. Multiple DNA damage-dependent and DNA damage-independent stress responses define the outcome of ATR/Chk1 targeting in medulloblastoma cells. *Cancer Lett.* **430**, 34–46 (2018).
210. Murga, M. et al. Exploiting oncogene-induced replicative stress for the selective killing of Myc-driven tumors. *Nat. Struct. Mol. Biol.* **18**, 1331–1335 (2011).
211. Roeschert, I. et al. Combined inhibition of Aurora-A and ATR kinases results in regression of MYCN-amplified neuroblastoma. *Nat. Cancer* **2**, 312–326 (2021).
212. Dammert, M. A. et al. MYC paralog-dependent apoptotic priming orchestrates a spectrum of vulnerabilities in small cell lung cancer. *Nat. Commun.* **10**, 3485 (2019).
213. Kelly, G. L. et al. Targeting of MCL-1 kills MYC-driven mouse and human lymphomas even when they bear mutations in p53. *Genes Dev.* **28**, 58–70 (2014).
214. Kretzner, L. et al. Combining histone deacetylase inhibitor vorinostat with aurora kinase inhibitors enhances lymphoma cell killing with repression of c-Myc, hTERT, and microRNA levels. *Cancer Res.* **71**, 3912–3920 (2011).
215. Pei, Y. et al. HDAC and PI3K antagonists cooperate to inhibit growth of MYC-driven medulloblastoma. *Cancer Cell* **29**, 311–323 (2016).
216. Lernoux, M. et al. Novel HDAC inhibitor MAKV-8 and imatinib synergistically kill chronic myeloid leukemia cells via inhibition of BCR-ABL/MYC-signaling: effect on imatinib resistance and stem cells. *Clin. Epigenet.* **12**, 69 (2020).
217. Zhao, X. et al. Disruption of the MYC-miRNA-EZH2 loop to suppress aggressive B-cell lymphoma survival and clonogenicity. *Leukemia* **27**, 2341–2350 (2013).
218. O'Donnell, K. A., Wentzel, E. A., Zeller, K. L., Dang, C. V. & Mendell, J. T. c-Myc-regulated microRNAs modulate E2F1 expression. *Nature* **435**, 839–843 (2005).
219. Dhanasekaran, R. et al. Anti-miR-17 therapy delays tumorigenesis in MYC-driven hepatocellular carcinoma (HCC). *Oncotarget* **9**, 5517–5528 (2018).
220. Lai, I. et al. Lipid nanoparticles that deliver IL-12 messenger RNA suppress tumorigenesis in MYC oncogene-driven hepatocellular carcinoma. *J. Immunother. Cancer* **6**, 125 (2018).
221. Pan, Y. et al. Synergistic inhibition of pancreatic cancer with anti-PD-L1 and c-Myc inhibitor JQ1. *Oncoimmunology* **8**, e1581529 (2019).
222. Devi, G. R. et al. In vivo bioavailability and pharmacokinetics of a c-MYC antisense phosphorodiamidate morpholino oligomer, AVI-4126, in solid tumors. *Clin. Cancer Res.* **11**, 3930–3938 (2005).
223. Tolcher, A. W. et al. Safety and activity of DCR-MYC, a first-in-class Dicer-substrate small interfering RNA (DsiRNA) targeting MYC, in a phase I study in patients with advanced solid tumors. *J. Clin. Oncol.* **33**, 11006–11006 (2015).
224. Villanueva, M. T. Long path to MYC inhibition approaches clinical trials. *Nat. Rev. Drug Discov.* <https://doi.org/10.1038/d41573-019-00055-2> (2019).
225. Shachaf, C. M. et al. Genomic and proteomic analysis reveals a threshold level of MYC required for tumor maintenance. *Cancer Res.* **68**, 5132–5142 (2008).
226. Jung, M. et al. A myc activity signature predicts poor clinical outcomes in myc-associated cancers. *Cancer Res.* **77**, 971–981 (2017).
227. Tran, D. et al. Functional genomics analysis reveals a myc signature associated with a poor clinical prognosis in liposarcomas. *Am. J. Pathol.* **185**, 717–728 (2015).

Acknowledgements

R.D. receives grant support from the NIH (grant CA222676). A.D. has received support from Lymphoma Research Foundation. A.S.H. has received support from Lundbreck Foundation. A.M.G. has received support from the NIH (grant CA196585). D.W.F. receives grant support from the NIH (grants CA208735, CA253180, CA188383 and CA184384).

Author contributions

R.D. and D.W.F. researched data for this manuscript. All authors contributed to the discussion of content and preparation of the manuscript.

Competing interests

D.W.F. is a consultant for Revolution Medicines, a company developing MYC pathway therapies, co-founder of Bachus and Molecular Decisions, and has had advisory roles for American Gene Technologies, Genon, Moderna and Regulus. The other authors declare no conflicts of interest.

Peer review information

Nature Reviews Clinical Oncology thanks M. Eilers, C.M. Eischen, E.V. Prochownik and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Supplementary information

The online version contains supplementary material available at <https://doi.org/10.1038/s41571-021-00549-2>.

© Springer Nature Limited 2021